

# Opportunities to Develop an Interagency Spatial Hierarchy for ESD Applications







## **Presentation Outline**

**Why develop an interagency hierarchy?**

**What is a spatial hierarchy – concepts and examples**

**Simple comparison of different national systems**

**Opportunities to formalize and map upper levels of the ESD hierarchy while revising and fully cross-walking systems**

**Example of cross-scale interactions and need for multi-scaled analysis and monitoring**

# Why develop an interagency hierarchy?

Policy - the 2008 MOU between the FS and NRCS for the NCSS states:

NRCS and FS mutually agree to complete a nationwide soil survey with ecological sites inventories by 2025;

FS and NRCS will be responsible for quality control for soil inventory.

# Why develop an interagency hierarchy?

Policy - the 2005 ESD MOU states:

The purpose of this MOU is to establish a Federal Interagency Team that will be responsible for developing a standardized method to be utilized by the BLM, FS, and NRCS to define, delineate and describe terrestrial ecological sites.

# Why develop an interagency hierarchy?

Practicality - ESD's need to be nested within relatively homogeneous climatic and physiographic zones.

These zones or broad-scale ecological units need to be adopted by all agencies for ESD's to be developed and coded consistently.

Corporate use of existing systems (FS national assessment examples - RPA, FIA, FHM, LANDFIRE) require maintaining existing systems.

Potential to revise where necessary then fully cross-walk systems to create an ESD hierarchy is real and needed.

# Spatial Hierarchy Concepts

Conditions and processes occurring across larger areas affect and often override those of smaller ecosystems, and the properties of smaller ecosystems emerge in the context of larger systems.

For example, a wetland embedded within a fire-prone landscape functions differently than one embedded within a fire-resistant landscape.


Moreover, environmental gradients affecting ecological patterns and processes change at different spatial scales, forming a natural spatial hierarchy.



# Important factors that interact to form ecosystems

- Climate
- Geology
- Physiography (slope, aspect, elevation)
- Soils
- Plants
- Animals
- Water
- Disturbance regimes

**The integration of multiple factors at relevant scales is all important in understanding ecosystems.**



# Spatial Hierarchy

At continental and regional scales, ecosystem patterns correspond with climatic regions, which change mainly due to latitudinal, orographic, and maritime influences.

Within climatic regions, landforms modify macroclimate, and affect the movement of organisms, the flow and orientation of watersheds, and the frequency and spatial pattern of disturbance by fire and wind.

Within climatic - geomorphic regions, water, plants, animals, soils, and topography interact to form ecosystems at more local scales



# Spatial Hierarchy

The challenge of ecosystem classification and mapping is to:

- Distinguish natural associations of ecological factors at relevant spatial scales
- Define ecological types or ESD's, and map ecological land units that reflect these different levels of organization
- Interpret the properties and dynamics of these systems for management.

# NRCS-BLM-FS Ecological Site Description Handbook Ecological Mapping Systems

<b>Hierarchical Planning and Analysis Levels</b>	<b>National Hierarchical Framework of Ecological Units<sup>1</sup></b>	<b>NRCS Soil Geography Hierarchy<sup>3</sup></b>	
<b>Continental and Region (Ecoregion)</b>	<b>Domain, Division, and Province (1:5,000,000-1:30,000,000)</b>	<b>Land Resource Region (LRR) (1:7,500,000), Climate zones</b>	
<b>Subregion</b>	<b>Section (1:3,500,000) and Subsection (1:250,000)</b>	<b>Major Land Resource Area (MLRA) (1:3,500,000) Land Resource Unit (LRU)/Common Resource Area (CRA) (1:1,000,000) General Soil Map (1:250,000)</b>	
<b>Landscape (watershed—5<sup>th</sup> unit of Hydrologic Unit Code)</b>	<b>Landtype Association (1:60,000)</b>	<b>Soil-geomorphic systems</b>	
<b>Land Unit (subwatershed—6<sup>th</sup> unit of Hydrologic Unit Code), grazing allotment, farm/ranch)</b>	<b>Landtype (1:24,000)</b>	<b>Detailed Soil Map (1:24,000)</b>	
	<b>Landtype Phase (1:12,000)</b>	<b>Soil Series (1:12,000)</b>	
<b>Individual Sites</b>	<b>Sampling plot</b>	<b>Soil Pedon</b>	

**From: Draft Interagency Ecological Site Handbook**

# Overview of climatic gradients






## Data Source

PRISM data (Parameter-elevation Regressions on Independent Slopes Model), developed by the Spatial Climate Analysis Service at Oregon State University.


Based on a 30 year period (1961-1990), and a 16 year period (1991-2007) for estimating recent shifts in climatic regimes.



**FreezeFreeDays**

Value





**MeanAnnualTemp**


**DEGREES\_F2**

- 24.4 - 30.7
- 30.8 - 33.6

<span style="display: inline-block; width: 15px; height: 15px; background-color: #e0e0e0; border: 1px solid black; margin-right: 5px;"></span> 33.7 - 36	<span style="display: inline-block; width: 15px; height: 15px; background-color: #4169e1; border: 1px solid black; margin-right: 5px;"></span> 41.8 - 43.9	<span style="display: inline-block; width: 15px; height: 15px; background-color: #32cd32; border: 1px solid black; margin-right: 5px;"></span> 51.2 - 53.6	<span style="display: inline-block; width: 15px; height: 15px; background-color: #ffff00; border: 1px solid black; margin-right: 5px;"></span> 61.3 - 63.5	<span style="display: inline-block; width: 15px; height: 15px; background-color: #8b4513; border: 1px solid black; margin-right: 5px;"></span> 71.2 - 73.4
<span style="display: inline-block; width: 15px; height: 15px; background-color: #d81b60; border: 1px solid black; margin-right: 5px;"></span> 36.1 - 37.9	<span style="display: inline-block; width: 15px; height: 15px; background-color: #4682b4; border: 1px solid black; margin-right: 5px;"></span> 44 - 46.2	<span style="display: inline-block; width: 15px; height: 15px; background-color: #32cd32; border: 1px solid black; margin-right: 5px;"></span> 53.7 - 56.1	<span style="display: inline-block; width: 15px; height: 15px; background-color: #ffa500; border: 1px solid black; margin-right: 5px;"></span> 63.6 - 65.8	<span style="display: inline-block; width: 15px; height: 15px; background-color: #800000; border: 1px solid black; margin-right: 5px;"></span> 73.5 - 77.9
<span style="display: inline-block; width: 15px; height: 15px; background-color: #800080; border: 1px solid black; margin-right: 5px;"></span> 38 - 39.7	<span style="display: inline-block; width: 15px; height: 15px; background-color: #00bfff; border: 1px solid black; margin-right: 5px;"></span> 46.3 - 48.6	<span style="display: inline-block; width: 15px; height: 15px; background-color: #7cfc00; border: 1px solid black; margin-right: 5px;"></span> 56.2 - 58.6	<span style="display: inline-block; width: 15px; height: 15px; background-color: #ff8c00; border: 1px solid black; margin-right: 5px;"></span> 65.9 - 68.5	
<span style="display: inline-block; width: 15px; height: 15px; background-color: #4169e1; border: 1px solid black; margin-right: 5px;"></span> 39.8 - 41.7	<span style="display: inline-block; width: 15px; height: 15px; background-color: #40e0d0; border: 1px solid black; margin-right: 5px;"></span> 48.7 - 51.1	<span style="display: inline-block; width: 15px; height: 15px; background-color: #adff2f; border: 1px solid black; margin-right: 5px;"></span> 58.7 - 61.2	<span style="display: inline-block; width: 15px; height: 15px; background-color: #cd853f; border: 1px solid black; margin-right: 5px;"></span> 68.6 - 71.1	




# USDA Zone Map




## Zone Avg. Annual Low

- 2  -40°F through -50°F
- 3  -30°F through -40°F
- 4  -20°F through -30°F
- 5  -10°F through -20°F
- 6  0°F through -10°F
- 7  10°F through 0°F
- 8  20°F through 10°F
- 9  30°F through 20°F
- 10  40°F through 30°F




**Mean Annual Precipitation**

PRECIP\_IN2







# Mean Annual Precipitation and FS Provinces





Mean Annual Precipitation	16.1 - 19.9	37 - 40.7	55.4 - 60.1
PRECIP_IN2	20 - 24.4	40.8 - 44.5	60.2 - 68.2
1.9 - 8.7	24.5 - 28.9	44.5 - 47.9	68.3 - 79.6
8.8 - 12.5	29 - 33.1	48 - 51.5	79.7 - 93.4
12.6 - 16	33.2 - 36.9	51.6 - 55.3	93.5 - 108.3












# Physiography, bedrock geology, surficial geology



# USGS Lithology 2010 (new data)



## Surficial


<all other values>


### UNIT\_NAME

Alluvial sediments, thick	Glaciofluvial ice-contact sediments, mostly sand and gravel, discontinuous
Alluvial sediments, thin	Glaciofluvial ice-contact sediments, mostly sand and gravel, thick
Basaltic and andesitic volcanic rocks	Glaciofluvial ice-contact sediments, mostly sand and gravel, thin
Calcareous biological sediments	Lacustrine sediments
Coastal zone sediments, mostly fine-grained	Organic-rich muck and peat, thick
Coastal zone sediments, mostly medium-grained	Organic-rich muck and peat, thin
Colluvial and alluvial sediments	Playa sediments
Colluvial sediments and loess	Proglacial sediments, mostly coarse-grained, discontinuous
Colluvial sediments and residual material	Proglacial sediments, mostly coarse-grained, thick
Colluvial sediments, discontinuous	Proglacial sediments, mostly coarse-grained, thin
Colluvial sediments, thin	Proglacial sediments, mostly fine grained, discontinuous
Eolian sediments on southern High Plains	Proglacial sediments, mostly fine grained, thick
Eolian sediments, mostly dune sand, thick	Proglacial sediments, mostly fine grained, thin
Eolian sediments, mostly dune sand, thin	Residual materials developed in alluvial sediments
Eolian sediments, mostly loess, thick	Residual materials developed in bedrock, discontinuous
Eolian sediments, mostly loess, thin	Residual materials developed in bedrock, thin
Glacial till sediments, mostly clayey, discontinuous	Residual materials developed in bedrock, with alluvial sediments, discontinuous
Glacial till sediments, mostly clayey, thick	Residual materials developed in bedrock, with alluvial sediments, thin
Glacial till sediments, mostly clayey, thin	Residual materials developed in carbonate rocks, discontinuous
Glacial till sediments, mostly sandy, discontinuous	Residual materials developed in carbonate rocks, thin
Glacial till sediments, mostly sandy, thin	Residual materials developed in fine-grained sedimentary rocks
Glacial till sediments, mostly silty, discontinuous	Residual materials developed in igneous and metamorphic rocks
Glacial till sediments, mostly silty, thick	Residual materials developed in sedimentary rocks, discontinuous
Glacial till sediments, mostly silty, thin	Residual materials developed in sedimentary rocks, thin
	Rhyolitic volcanic rocks
	Water




# USGS Lithology and NRCS LRR's



 mlra\_v42\_LRR

# USGS Lithology and FS Sections







# USGS Lithology and FS Subsections






# Ssurgo water holding capacity and FS Subsections





# Subsections and Historical Vegetation






Three distinct, mappable landscape ecosystems (LTA's)  
with different fire regimes, habitat quality, etc.

Fire-resistant moist-mesic sugar maple – basswood PNV  
On loamy morainal ecosystems

Moderately fire-prone dry-mesic red – white pine PNV  
On loamy sand ice-contact ecosystems

Highly fire-prone xeric jackpine PNV  
On coarse sandy outwash ecosystems





# **Hierarchical Mapping Systems**

**USDA NRCS LRR's, MLRA's, Statsgo, Ssurgo**

**USDA FS National Hierarchy of Ecological Units**

**EPA Ecoregions**

**NatureServe**

**USGS**

# NRCS Land Resource Regions



## Land Resources Regions

mra\_v42


<all other values>

LRR\_NAME

- |                                                                |                                                         |                                                        |                                                                        |
|----------------------------------------------------------------|---------------------------------------------------------|--------------------------------------------------------|------------------------------------------------------------------------|
| Alberta, Alaska                                                | Central Great Plains Winter Wheat and Range Region      | Northeastern Forage and Forest Region                  | Rocky Mountain Range and Forest Region                                 |
| Atlantic and Gulf Coast Lowland Forest and Crop Region         | Eastern Central Farming and Forest Region               | Northern Alaska                                        | South Atlantic and Gulf Slope Cash Crops, Forest, and Livestock Region |
| California Subtropical Fruit, Truck, and Specialty Crop Region | Florida Subtropical Fruit, Truck Crop, and Range Region | Northern Atlantic Slope Diversified Farming Region     | Southern Alaska                                                        |
| Caribbean Region                                               | Hawaii Region                                           | Northern Great Plains Spring Wheat Region              | Southwest Plateaus and Plains Range and Cotton Region                  |
| Central Feed Grains and Livestock Region                       | Interior Alaska                                         | Northern Lake States Forest and Forage Region          | Southwestern Prairies Cotton and Forage Region                         |
| Mississippi Delta Cotton and Feed Grains Region                | Lake State Fruit, Truck Crop, and Dairy Region          | Northwestern Forest, Forage, and Specialty Crop Region | Western Alaska                                                         |
|                                                                | Mississippi Delta Cotton and Feed Grains Region         | Northwestern Wheat and Range Region                    | Western Great Plains Range and Irrigated Region                        |
|                                                                |                                                         | Pacific Basin Region                                   | Western Range and Irrigated Region                                     |



# NRCS LRR's and FS Provinces



## Land Resources Regions


mra\_v42

<all other values>

LRR\_NAME

- |                                                                |                                                         |                                                        |                                                               |
|----------------------------------------------------------------|---------------------------------------------------------|--------------------------------------------------------|---------------------------------------------------------------|
| Alberta Alaska                                                 | Central Great Plains Winter Wheat and Range Region      | Northeastern Forage and Forest Region                  | Rocky Mountain Range and Forest Region                        |
| Atlantic and Gulf Coast Lowland Forest and Crop Region         | Eastern Central Farming and Forest Region               | Northern Alaska                                        | South Atlantic Slope Cash Crops, Forest, and Livestock Region |
| California Subtropical Fruit, Truck, and Specialty Crop Region | Florida Subtropical Fruit, Truck Crop, and Range Region | Northern Atlantic Slope Diversified Farming Region     | Southern Alaska                                               |
| Caribbean Region                                               | Hawaii                                                  | Northern Great Plains Spring Wheat Region              | Southwest Plateaus and Plains Range and Cotton Region         |
| Central Feed Grains and Livestock Region                       | Hawaii Region                                           | Northern Lake States Forest and Forage Region          | Southwestern Plateaus Cotton and Forage Region                |
|                                                                | Interior Alaska                                         | Northwestern Forest, Forage, and Specialty Crop Region | Western Alaska                                                |
|                                                                | Lake State Fruit, Truck Crop, and Dairy Region          | Northwestern Wheat and Range Region                    | Western Great Plains Range and Irrigated Region               |
|                                                                | Mississippi Delta Cotton and Feed Grains Region         | Pacific Basin Region                                   | Western Range and Irrigated Region                            |

# NRCS LRR's and FS Sections



## Land Resources Regions


mra\_v42

<all other values>


LRR\_NAME

Arctic Alaska	Central Great Plains Winter Wheat and Range Region	Northeastern Forage and Forest Region	Rocky Mountain Range and Forest Region
Atlantic and Gulf Coast Lowland Forest and Crop Region	Eastern Central Farming and Forest Region	Northern Alaska	South Atlantic and Gulf Slope Cash Crops, Forest, and Livestock Region
California Subtropical Fruit, Truck, and Specialty Crop Region	Florida Subtropical Fruit, Truck Crop, and Range Region	Northern Atlantic Slope Diversified Farming Region	Southern Alaska
Caribbean Region	Hawaii	Northern Great Plains Spring Wheat Region	Southwest Plateaus and Plains Range and Cotton Region
Central Feed Grains and Livestock Region	Hawaii Region	Northern Lake States Forest and Forage Region	Southwestern Plateaus Cotton and Forage Region
	Interior Alaska	Northwestern Forest, Forage, and Specialty Crop Region	Western Alaska
	Lake State Fruit, Truck Crop, and Dairy Region	Northwestern Wheat and Range Region	Western Great Plains Range and Irrigated Region
	Mississippi Delta Cotton and Feed Grains Region	Pacific Basin Region	Western Range and Irrigated Region

# NRCS MLRA's and FS Sections




## NRCS MLRA's and FS Subsections






# EPA Level 3 Ecoregions and FS Sections



# EPA Level 3 Ecoregions and FS Subsections






### A New Map of Standardized Terrestrial Ecosystems of the Conterminous United States



Professional Paper 1768

U.S. Department of the Interior  
U.S. Geological Survey





“A New Map of Standardized Terrestrial Ecosystems of the Conterminous United States”  
(USGS -NatureServe 2009) and Section Boundaries

The boundaries of different mapping systems delineating broader-scale ecological regions are converging, most likely due to improved technology.

The principal differences are interpretations of scale relationships.


The opportunity to develop an interagency hierarchy for use in ESD applications, while revising respective agency complementary systems, has never been more possible.

The barriers are not scientific, they are institutional.


# **Example of the need for use of a spatial hierarchy**

**While conducting multi-scaled analysis and monitoring, and**

**Evaluating cross-scale interactions**







## Thresholds of Climate Change in Ecosystems

**U.S. Climate Change Science Program**  
Synthesis and Assessment Product 4.2

January 2009

5.2.1 Role of Monitoring: “Because climate change effects are likely to interact with patterns and processes across spatial and temporal scales, it is clear the monitoring strategies must be integrated across scales.”



## Thresholds of Climate Change in Ecosystems

**U.S. Climate Change Science Program**  
Synthesis and Assessment Product 4.2

January 2009

“First and foremost, the earth’s surface must be hierarchically stratified (for example, using the MLRA’s and Ecological Site Description System of the U.S. D.A. National Resources Conservation Service and U.S. Forest Service ecoregions), and conceptual or simulation models of possible impacts and feedbacks must be specified for each stratum (Herrick et al., 2006).



## Thresholds of Climate Change in Ecosystems

U.S. Climate Change Science Program  
Synthesis and Assessment Product 4.2

“The models are used to develop scenarios and to identify key properties and processes that are likely to be associated with abrupt changes.

Second, simultaneous multiple-scale monitoring should be implemented at up to three spatial scales based on these scenarios and the recognition of pattern-and-process coupling developed in the models (Bestelmeyer, 2006), which may feature cross-scale interactions (Peters et al., 2004).”




# The Colorado Plateau Snowpack – Dust Interaction

Dust originating from larger surrounding shrubland and grassland dominated landscapes is being deposited within alpine zones in Colorado.

This has caused snowpacks to melt 35 – 45 days earlier than normal, affecting hydrologic function, urban water supply, and recreation (skiing).



# The Colorado Plateau Snowpack – Dust Interaction



It may also be affecting the phenology of plants, movement of tree lines, the synchrony of pollinators and flowering plants, seasonality of soil temperature and moisture regimes, and a host of other processes.



# The Colorado Plateau Snowpack – Dust Interaction

This raises several questions related to:

- Causal relationships
- Scale of observation for monitoring and detection
- Interactions which may be occurring across scales



# Causal Relationships



Dust is originating from drier, lower lying areas where destabilization of soil crusts, loss of vegetative cover, and high winds facilitate higher elevation deposition.

The effect of regional sources of dust vary at a landscape scale, with altered albedo in alpine areas differing from lower elevation forests areas.



# Causal Relationships


The questions are, is desertification taking place because of:

- (i) recent drought or climate change?
- (ii) anthropogenic forcing via land-use?
- (iii) a natural range of variability phenomenon?
- (iv) interactions of the above?


# Recent shifts in climatic regimes


Comparison of components of climatic regimes of the 1961-1990 versus 1991-2007 periods







### Change\_Winter\_Temp






**Change\_Summer\_Temp**







### Change winter precipitation









### Change growing season precipitation





### Percent Change Growing Season Precipitation



# Interactions that may be occurring across scales


It is possible that climate change is the driving force

It is possible that land use is the driving force

It is possible that long-term natural variability in climate is the driving force



**Fig. 5** Climatic shift years in the normalized (1725–1999) winter PNAI series, identified using a two-sample  $t$ -test between the first and second half of 30-year moving windows. Significance levels are indicated by *full* ( $P < 0.01$ ) and *dashed* ( $P < 0.05$ ) horizontal lines. Climatic shift years (vertical dashed lines) were defined as the years with highest absolute  $t$ -value. Years of climatic shifts identified in other proxy records of Pacific climate variability and solar activity (Crowley 2000) are also indicated. *Filled symbols* represent climatic shifts within a 3 year range from the shifts in winter PNAI



Multi-century variability in the Pacific North American circulation pattern reconstructed from tree rings. Trouet and Taylor. *Clim Dyn* (2010) 35:953–963.

“Positive PNA phases produce below average snow accumulation in western North America as a result of warm temperatures and decreased precipitation.”

# Interactions that may be occurring across scales

It is also possible that implementation of best management practices has been effective in reducing effects of grazing, recreation, or other anthropogenic impacts.

And that the rate of desertification due to climate change or a positive Pacific North American circulation pattern has been slowed through these practices.



# Interactions that may be occurring across scales

However, monitoring meso-scale trends in land use in the absence of macro-scale monitoring might lead to conclusions that BMP's are ineffective, that opportunity costs of limiting resource use are being incurred, and BMP's should be adjusted accordingly.



# Interactions that may be occurring across scales

Similarly, assessing landscape level conditions and processes, and implementing adaptation strategies at the landscape scale may not be effective without implementing adaptation strategies at the meso-scale.



# Scale of observation for monitoring and detection

Macro-scale – monitoring climate change

Meso-scale – monitoring land use, BMP's

Landscape scale – monitoring snow, dust, vegetation within Alpine zones

Local scale – monitoring response of various species, hydrology, other phenomena of interest



## Cross-scale interactions

The bottom line is broader scale stressors may override finer-scale conditions and actions that are effective at reducing adverse cumulative effects.

And finer-scale processes (destabilized soil crusts, coalescence of open patches) may propagate upward through the system to alter broader scale patterns (dust production, snowpack melt).



# Multi-scaled Monitoring

Designing inventory and monitoring programs that employ concepts of hierarchical structures is therefore needed for assessing climate change as well as other stressors.

We have that opportunity via the interagency ESD effort.